Table D10. Individual hydrocarbon and total petroleum hydrocarbon concentrations (in μ g/g wet wt.) for ribbed mussels taken from Mill Creek marsh, a reference site. 1-3

Sample ID	Nonane (n-C ₉)	Decane (n-C ₁₀)	Undecane (n-C ₁₁)	Dodecane (n-C ₁₂)	Tridecane (n-C ₁₃)	Tetradecane (n-C ₁₄)	Pentadecane (n-C ₁₅)	Hexadecane (n-C ₁₆)	Heptadecane (n-C ₁₇)	Pristane	Octadecane (n-C ₁₈)	Phytane	Nonadecane (n-C ₁₉)	Eicosane (n-C ₂₀)	Heneicosane (n- C_{21})	Docosane (n-C ₂₂)	Tricosane (n-C ₂₃)
First Collection																	
197020407	0.98	nd	nd	nd	nd	nd	0.14	0.13	0.29	0.14	nd	nd	0.24	nd	2.41	0.81	nd
197020417	1.28	nd	nd	nd	nd	nd	nd	nd	0.12	nd	nd	nd	0.35	nd	1.52	1.03	nd
197020418	0.61	nd	nd	nd	nd	nd	nd	0.15	0.26	nd	nd	nd	0.13	nd	1.90	0.76	nd
197020419	0.42	nd	nd	nd	nd	nd	0.12	nd	0.19	nd	nd	nd	nd	nd	0.69	1.78	nd
197020420	nd	nd	nd	nd	nd	nd	nd	0.10	0.11	nd	nd	nd	nd	nd	1.04	nd	nd
Average ⁴	0.68	nd	nd	nd	nd	nd	< MDL	0.09	0.19	< MDL	nd	nd	0.16	nd	1.51	0.95	nd
Std Dev	0.46	-	-	-	-	-	-	0.05	0.08	-	-	-	0.13	-	0.68	0.53	-
Second Collection																	
497051417	0.34	nd	nd	nd	nd	nd	nd	nd	0.76	nd	nd	0.10	nd	nd	1.31	nd	nd
497051418	0.42	nd	nd	nd	nd	nd	nd	nd	0.47	nd	nd	nd	nd	nd	1.18	nd	nd
497051419	0.27	nd	nd	nd	nd	nd	nd	nd	1.06	nd	nd	0.11	0.15	nd	3.43	1.04	4.22
497051420	nd	nd	nd	nd	nd	nd	nd	nd	1.29	nd	nd	nd	nd	nd	2.27	nd	nd
497051421	0.68	nd	nd	nd	nd	nd	nd	nd	1.15	nd	0.16	0.12	1.44	nd	1.85	nd	nd
Average ⁴	0.37	nd	nd	nd	nd	nd	nd	nd	0.95	nd	< MDL	< MDL	0.35	nd	2.01	< MDL	< MDL
Std Dev	0.21	-	-	-	-	-	-	-	0.33	-	-	-	0.61	-	0.91	-	-
MDL	0.24	0.06	0.12	0.12	0.11	0.11	0.11	0.09	0.08	0.10	0.10	0.10	0.11	0.21	0.38	0.68	2.47

Table D10. Continued. 1-3

Sample ID	Tetracosane (n-C ₂₄)	Pentacosane (n- C_{25})	Hexacosane (n-C ₂₆)	Heptacosane (n-C ₂₇)	Octacosane (n-C ₂₈)	Nonacosane (n-C ₂₉)	Triacontane (n-C ₃₀)	n-Hentriacontane (n-C ₃₁)	Dotriacontane (n-C ₃₂)	Tritriacontane $(n-C_{33})$	Tetratriacontane $(n-C_{34})$	Pentatriacontane (n-C ₃₅)	Hexatriacontane $(n-C_{36})$	Heptatriacontane (n-C ₃₇)	Octatriacontane (n-C ₃₈)	Nonatriacontane (n-C ₃₉)	Tetracontane (n-C ₄₀)
First Collection																	
197020407	nd	0.30	0.19	nd	nd	nd	nd	0.48	0.64	0.68	0.75	0.83	0.77	0.80	0.56	0.45	0.26
197020417	nd	nd	nd	0.08	nd	nd	nd	0.52	0.56	0.58	0.59	0.63	0.57	0.53	0.43	0.31	0.18
197020418	nd	nd	nd	nd	nd	nd	nd	0.23	0.36	0.42	0.44	0.50	0.49	0.44	0.40	0.28	0.17
197020419	nd	nd	nd	nd	nd	nd	nd	0.16	0.24	0.26	0.27	0.32	0.32	0.32	0.25	0.18	0.11
197020420	nd	nd	nd	nd	nd	nd	nd	56.6	5.12	2.11	0.22	0.25	0.24	0.26	0.19	0.14	0.09
Average⁴	nd	< MDL	< MDL	< MDL	nd	nd	nd	11.6	1.38	0.81	0.45	0.51	0.48	0.47	0.37	0.27	0.16
Std Dev	-	-	-	-	-	-	-	25.1	2.09	0.74	0.22	0.23	0.21	0.21	0.15	0.12	0.07
Second Collection																	
497051417	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
497051418	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	0.09	nd	nd
497051419	nd	nd	nd	nd	nd	nd	nd	0.25	1.36	0.91	nd	nd	nd	nd	0.08	nd	nd
497051420	nd	nd	nd	nd	nd	nd	nd	0.84	4.89	2.09	nd	nd	nd	nd	0.10	0.06	nd
497051421	nd	nd	nd	nd	nd	nd	nd	0.59	1.81	1.52	nd	nd	nd	nd	0.08	nd	nd
Average⁴	nd	nd	nd	nd	nd	nd	nd	0.36	1.64	0.93	nd	nd	nd	nd	< MDL	< MDL	nd
Std Dev	-	-	-	-	-	-	-	0.35	1.97	0.90	-	-	-	-	-	-	-
MDL	0.29	0.27	0.11	0.08	0.20	0.56	0.48	0.11	0.14	0.11	0.09	0.10	0.10	0.18	0.08	0.06	0.06

Table D10. Continued. 1-3

Sample ID	Total Petroleum Hydrocarbons ⁵	Total Concentrations of Individual Hydrocarbons ^{67,15}	Total: Pristane + Phytane ^{6,15}	Pristane/n-C-7 ¹⁶	Phytane/n-C₁፥¹6	Pristane/Phytane ¹⁶	Total: Odd No Carbons ^{6,8,15}	Total: Even No Carbons ^{6,9,15}	Carbon Preference Index (CPI) ^{10,16}	Sum: C ₁₀ -C ₁₂ -C ₁₄ ^{6,11,15}	Sum: C ₂₂ -C ₂₄ -C ₂₈ -C ₂₈ 61215	Weathering Index (WI) ^{13,16}
First Collection												
197020407	148	14.4	nd	0.48	_	_	9.28	4.90	1.89	nd	nd	_
197020417	99.9	12.1	nd	-	_	_	7.72	4.24	1.82	nd	1.33	_
197020418	111	10.3	nd	_	_	_	6.62	3.62	1.83	nd	nd	_
197020419	64.6	8.46	nd	_	-	_	nd	3.86	-	nd	2.08	-
197020420	141	69.7	nd	-	-	-	62.5	7.14	8.76	nd	nd	-
Average ⁴	113	23.0	nd	-	-	-	18.1	4.75	3.81 ¹⁷	nd	<mdl< td=""><td>_</td></mdl<>	_
Std Dev	33.6	26.2	-	-	-	-	24.9	1.42	-	-	-	-
Second Collection												
497051417	110	nd	nd	-	-	-	nd	nd	-	nd	nd	-
497051418	90.0	nd	nd	-	-	-	nd	nd	-	nd	nd	-
497051419	208	14.7	nd	-	-	-	11.1	3.48	3.18	nd	1.34	-
497051420	233	15.2	nd	-	-	-	8.72	6.33	1.38	nd	nd	-
497051421	167	12.8	nd	-	0.78	-	9.26	3.35	2.76	nd	nd	-
Average ⁴	162	10.9	nd	-	_	-	7.59	3.23	2.35 ¹⁷	nd	< MDL	_
Std Dev	61.4	4.58	-	-	-	-	3.01	1.99	-	-	-	-
MDL	53.6	8.19 ¹⁴	0.19 ¹⁴				5.09 ¹⁴	2.91 ¹⁴		0.29 ¹⁴	1.29 ¹⁴	

Table D10. Continued.

Footnotes:

- 1 The concentrations of the individual aliphatic hydrocarbons and the total petroleum hydrocarbons were determined using external standard calculations.
- When an individual aliphatic hydrocarbon was not detected, its concentration was replaced by nd.
- The concentrations for n-C₈ will be not reported, since it was difficult to identify this peak in samples and to determine MDL for n-C₈.
- If all concentrations are nd, the average is replaced with nd. When there is at least one number in the data set to be averaged, each nd is replaced with 1/2*MDL, and an average is calculated. If this numeric value is less than the MDL, the average is replaced by < MDL; otherwise, the average is the calculated value. When a numeric value is found for the average, the standard deviation is then determined using the same number set used to calculate the average.
- Determined from the total peak areas in the chromatogram from n-C₈ to n-C₄₀ minus any contributions from the internal standard areas.
- These formulae use 1/2MDL values for each analyte not detected.
- Sum of the concentrations of the individual aliphatic hydrocarbons $n-C_9$ through $n-C_{40}$ plus the concentrations of pristane and phytane.
- The total of the concentrations of the aliphatic hydrocarbons with an odd number of carbon atoms.
- The total of the concentrations of the aliphatic hydrocarbons with an even number of carbon atoms. The contribution of n-C₈ is not included in the total.
- Carbon Preference Index (CPI) is defined as the ratio of the total of the concentrations of the aliphatic hydrocarbons with an odd number of carbons to the total concentration of the alphatic hydrocarbons with an even carbon number.
- 11 The total of the concentrations of n-C₁₀, n-C₁₂, and n-C₁₄.
- The total of the concentrations of n-C₂₂, n-C₂₄, n-C₂₆, and n-C₂₈.
- Weathering Index (WI) is defined as the ratio of the total concentration of n-C₁₀, n-C₁₂, and n-C₁₄ to the total concentration of n-C₂₂, n-C₂₄, n-C₂₆, and n-C₂₈.
- These MDL values are calculated with the same summation formulae as the samples using the individual hydrocarbon MDL values.
- The summation totals for the samples are compared with calculated MDL values obtained using the same summation formulae as the samples. When these sample totals were less than the total MDL, its value was replaced by nd. The averages and standard deviations for the totals were treated in the same way as the individual hydrocarbons; see footnote 4.
- Numerical values of the CPI, WI, and the ratios: pristane/n-C₁₇, phytane/n-C₁₈, and pristane/phytane, will be calculated only when the defined quantity for each index or ratio has a numeric value.
- 17 These results are not true averages, instead they are the ratios of the averages of the defined quantities, if these averages exist.